

# INSTRUCTIONS FOR THE PHYSICIAN

## HOW TO PREPARE THE GONAPEPTYL® DEPOT SUSPENSION:

Since successful treatment depends on correct preparation of the suspension, the following instructions must be strictly followed.

### 1. Preparations:

- Take the package of Gonapeptyl® Depot out of the refrigerator.
- Remove the cap from the disposable syringe containing the controlled-release microcapsules.
- Open the package containing the connector **without** removing the connector.
- Screw the syringe containing the controlled-release microcapsules onto the connector in the package, then remove it.
- Screw the syringe containing the solvent tightly on the free end of the connector and ensure that it fits tightly (1).




Fig. 1

### 2. Reconstitution of a suspension:

- Empty the liquid into the syringe containing the powder (2), then push it back and forth into the first syringe. The first two or three times **without pushing the injection rod all the way in**. Repeat this about 10 times or until you have a homogeneous milky-like suspension.
- Some foam may be created during preparation of the suspension. It is essential that the foam be dissolved or removed from the syringe before giving the injection.
- The liquid has to be transferred between syringes until a homogeneous milky-like suspension is achieved (3).




Fig. 2: Mixing




Fig. 3: Mix abt. 10 times

### 3. Injection:

- Remove the connector together with the empty syringe.
- Mount the injection needle on the syringe with the ready-to-use suspension.
- Inject subcutaneously or deep into the muscle **immediately**. (Within 3 minutes)

packed by

**FERRING**  
PHARMACEUTICALS

St. Prex, Switzerland

# AANWYSINGS AAN DIE GENEESHEER OM GONAPEPTYL® DEPOT SUSPENSIE VOOR TE BEREI:

Aangesien suksesvolle behandeling afhang van die korrekte voorbereiding van die suspensie, moet die volgende instruksies streng gevolg word:

## 1. Voorbereiding:

- Verwyder die pak met Gonapeptyl® Depot uit die yskas.
- Verwyder die doppie van die weggooibare inspuiting wat die vrystelling-beheerde mikrokapsules bevat.
- Maak die pakkie wat die verbindstuk bevat oop **sonder** om die verbindstuk te verwyder.
- Skroef die glasspuit wat die vrystelling-beheerde mikrokapsules bevat aan die verbindstuk vas en verwyder dit dan.
- Skroef die glasspuit, wat die oplosmiddel bevat, styf aan die vry kant van die verbindstuk vas en maak seker dat dit styf pas (1).




Fig. 1

## 2. Aanmaak van die suspensie:

- Spuit die vloeistof in die glasspuit met die poeier in (2) en druk dit heen en weer in die eerste glasspuit in. Die eerste twee of drie keer **sonder om die spuitstang al die pad in te druk**. Herhaal die proses omtrent 10 keer of totdat 'n homogene, melkerige suspensie verkry is (3).
- 'n Bietjie skuim mag vorm gedurende die voorbereidingsproses van die suspensie. Dit is noodsaaklik dat die skuim opgelos of verwyder moet word van die glasspuit voor toediening.




Fig. 2: Vermenging




Fig. 3: Meng ongeveer 10 keer

## 3. Toediening:

- Verwyder beide die verbindstuk en die leë glasspuit.
- Verbind die spuitnaald met die glasspuit met die gereed-gebruik suspensie.
- Dien **dadellik** subkutaan of diep intramuskulêr toe. (Toediening moet binne 3 minute vanaf voorbereiding geskied).

Gepak deur

**FERRING**  
PHARMACEUTICALS

St. Prex, Switzerland