

CONTRA-INDICATIONS:

Hypersensitivity to the ingredients.

WARNINGS AND SPECIAL PRECAUTIONS:

Contains sucrose: Patients with the rare hereditary condition such as fructose intolerance, glucose-galactose mal-absorption or sucrose-isomaltase insufficiency should not take Vita-Thion.

Contains lactose monohydrate: Patients with the rare hereditary condition of galactose intolerance e.g. galactosaemia, Lapp lactase deficiency, glucose-galactose mal-absorption or fructose intolerance should not take Vita-Thion.

DOSAGE AND DIRECTIONS FOR USE:

Adults: 1 to 2 sachets dissolved in water at breakfast.

Children: 2 to 5 years: half a sachet at breakfast.

5 to 15 years: half a sachet at breakfast and half a sachet at lunch.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

Treatment is symptomatic.

IDENTIFICATION:

Pink to red and white granules.

PRESENTATION:

Boxes containing 2, 10, 20, 30 and 60 sachets.

STORAGE INSTRUCTIONS:

Store at or below 25 °C in a dry place.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER:

M/33/250

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

Adcock Ingram Limited

1 New Road

Erand Gardens

Midrand, 1685

Private Bag X69

Bryanston, 2021

www.adcock.com

Under license of LES LABORATOIRES SERVIER

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

October 1999