

### 1.3.1.1 PROFESSIONAL INFORMATION FOR MEDICINES FOR HUMAN USE

#### SCHEDULING STATUS

**S3**

#### PROPRIETARY NAME AND DOSAGE FORM

**MENIVERT 24 mg TABLETS**

#### COMPOSITION

Each tablet of MENIVERT 24 mg TABLETS contains 24 mg of betahistine dihydrochloride.

##### *Excipients:*

Citric acid anhydrous, crospovidone, hydrogenated vegetable oil, lactose monohydrate, maize starch, microcrystalline cellulose, povidone

Contains sugar: Lactose monohydrate 150 mg

#### CATEGORY AND CLASS

A 5.6 Histamine

#### PHARMACOLOGICAL ACTION

##### **Pharmacodynamic properties**

Betahistine is an analogue of histamine. It improves the microcirculation of the labyrinth resulting in reduced endolymphatic pressure. Pharmacological testing in animals has shown that the blood circulation in the *striae vascularis* of the inner ear improves, probably by means of a relaxation of the precapillary sphincters of the microcirculation of the inner ear.

In pharmacological studies, betahistine was found to have weak H<sub>1</sub> receptor agonistic and considerable H<sub>3</sub> antagonistic properties in the central nervous system and the autonomic nervous system.

Betahistine was also found to have a dose dependant inhibiting effect on spike generation of neurons in lateral and medial vestibular nuclei.

### **Pharmacokinetic properties**

#### **Absorption**

Following oral administration betahistine is well absorbed.

#### **Distribution**

There is little or no binding to plasma proteins.

#### **Metabolism**

Betahistine undergoes hepatic biotransformation, with evidence for the formation of 2-pyridylacetaldehyde and 2-(2-aminoethyl)pyridine.

#### **Elimination**

It is excreted in urine as the metabolite 2-pyridylacetic acid within 24 hours. No unchanged betahistine has been detected.

Betahistine is excreted in breast milk at approximately the same level as found in plasma.

### **INDICATIONS**

MENIVERT 24 mg TABLETS is indicated for the symptomatic treatment of the vertigo

associated with Ménière's syndrome.

## **CONTRAINDICATIONS**

MENIVERT 24 mg TABLETS is contraindicated in:

- Patients with hypersensitivity to betahistine or to any of the ingredients contained in MENIVERT 24 mg TABLETS (see COMPOSITION).
- Patients with pheochromocytoma. As betahistine is a synthetic analogue of histamine it may induce the release of catecholamines from the tumour resulting in severe hypertension.

## **WARNINGS AND SPECIAL PRECAUTIONS**

Caution is advised in the treatment of patients with peptic ulcer or a history of peptic ulceration, because of the occasional dyspepsia encountered in patients on MENIVERT 24 mg TABLETS.

Patients with bronchial asthma should be monitored carefully during the treatment with MENIVERT 24 mg TABLETS.

Caution is advised in prescribing MENIVERT 24 mg TABLETS to patients with either urticaria, rashes or allergic rhinitis, because of the possibility of aggravating these symptoms.

Caution is advised in patients with severe hypotension.

Caution is advised in patients with porphyria.

If MENIVERT 24 mg TABLETS are to be administered subsequent to the treatment with an antihistamine and this treatment is stopped abruptly, withdrawal symptoms such as sleep disorders and agitation could appear because of the sedative action of antihistamines. Treatment with the antihistamine should be tapered over approximately six days.

#### *Effects on ability to drive and use machines*

Since adverse reactions such as drowsiness have been reported in patients receiving MENIVERT 24 mg TABLETS, patients should not drive, use machinery or perform any tasks that require concentration, until they are certain that MENIVERT 24 mg TABLETS does not adversely affect their ability to do so (see SIDE EFFECTS).

#### *Excipients*

##### *Lactose warning:*

MENIVERT 24 mg TABLETS contain lactose which may have an effect on the glycaemic control of patients with diabetes mellitus. Patients with rare hereditary problems of galactose intolerance e.g. galactosaemia, the Lapp lactase deficiency or glucose-galactose malabsorption should not take MENIVERT 24 mg TABLETS.

## **INTERACTIONS**

There are no studies on the interactions with the following medicines: vasodilators, psychotropic medicines, in particular sedatives, tranquillisers and neuroleptics, parasympatholytics, and vitamins.

MENIVERT 24 mg TABLETS should not be concomitantly administered with antihistamines.

MENIVERT 24 mg TABLETS metabolism may be inhibited by medications that inhibit

monoamine-oxidase (MAO) including MAO subtype B (e.g. selegiline). Caution is recommended when using MENIVERT 24 mg TABLETS and MAO inhibitors (including MAO-B selective) concomitantly.

### **HUMAN REPRODUCTION**

The safety of MENIVERT 24 mg TABLETS during pregnancy and lactation has not been established.

### **DOSAGE AND DIRECTIONS FOR USE**

The usual daily dose is 24 mg to 48 mg of MENIVERT 24 mg TABLETS in divided doses. One 24 mg tablet can be administered twice daily.

The dosage should be individually adapted according to the response.

MENIVERT 24 mg TABLETS are not recommended for use in children and adolescents below the age of 18 years due to a lack of data on safety and efficacy.

### **SIDE EFFECTS**

MENIVERT 24 mg TABLETS can cause the following side effects:

#### **Immune system disorders**

*Frequency unknown:* Hypersensitivity reactions, e.g. anaphylaxis

#### **Nervous system disorders**

*Frequent:* Headache

*Less frequent:* Head pressure

*Frequency unknown:* Occasional drowsiness

**Cardiac disorders**

*Less frequent:* Palpitations, tightness of the chest

**Respiratory, thoracic and mediastinal disorders**

*Less frequent:* Exacerbation of pre-existing bronchial asthma

**Gastrointestinal disorders**

*Frequent:* Nausea, dyspepsia

*Less frequent:* Retching, heartburn, gastric discomfort and pain, flatulence

Gastric disorders can normally be avoided by taking MENIVERT 24 mg TABLETS with or after a meal, or by reducing the dosage.

**Skin and subcutaneous tissue disorders**

*Frequency unknown:* Cutaneous and subcutaneous hypersensitivity reactions, in particular angioneurotic oedema, urticaria rash, and pruritus

**KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENTS****Symptoms**

In case of overdose, the following symptoms, analogous to histamine overdose, might occur: headache, redness of the face, vertigo, tachycardia, hypotension, bronchial spasm, oedema, in particular oedema of the mucosa of the upper respiratory tract (Quincke's oedema).

**Treatment**

There is no specific antidote to MENIVERT 24 mg TABLETS.

**IDENTIFICATION**

MENIVERT 24 mg TABLETS are white, round, flat tablets with bevelled edges, breakline on one side. The tablets can be divided into equal halves.

**PRESENTATION**

20 tablets are packed in a polyvinylchloride, polyethylene, polyvinylidene chloride film sealed with aluminium foil backing. There are 10 tablets per blister strip and two blister strips are packed into an outer cardboard carton.

**STORAGE INSTRUCTIONS**

Store at or below 25 °C.

Keep in original packaging until required for use.

**KEEP OUT OF REACH OF CHILDREN.**

**REGISTRATION NUMBER**

46/5.6/0960

**NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF  
REGISTRATION**

PHARMACARE LIMITED

Healthcare Park

Woodlands Drive

Woodmead 2191

**DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION FOR MEDICINES  
FOR HUMAN USE**

Date of registration: 02 June 2017

Date of the most recent amendment to the professional information as approved by the

Authority: 02 June 2017

ZA\_MENITAB\_1706\_01